

# Trends and Factors Impacting Healthcare Charges and Length of Stay for Cholecystectomies: A New York State Population-based Analysis

Aria Darbandi,<sup>1</sup> Christina Chopra.<sup>2</sup>

## Abstract

**Background:** Gallbladder disease confers a significant economic toll on the United States healthcare system. This study aims to characterize current trends and features of the cholecystectomy population and identify factors that influence the length of stay and total charges. **Methods:** Case information was extracted for laparoscopic and open cholecystectomies from 2013-2016 using the New York Statewide Planning and Research Cooperative System (SPARCS) database. Descriptive, comparative, and multivariable linear regression analysis was conducted on 58,141 cases assessing age group, race, gender, admission presentation, surgical technique, insurance status, year of operation and severity of illness by the length of stay and total charges. **Results:** Of all procedures, 91.6% were laparoscopic, and 79.4% were emergent on admission. Total procedures trended down, while laparoscopic and emergent cases steadily increased ( $p < 0.0001$ ). Total charges increased during the study period, while the length of stay decreased ( $p < 0.0001$ ). Open and emergent procedures were associated with a higher cost and longer inpatient stays ( $p < 0.0001$ ). Open procedures were proportionally more common among elderly, male patients, and in elective cases ( $p < 0.0001$ ). Emergent presentation was more common in females, non-whites, and younger patients ( $p < 0.0001$ ). Regression model showed that male gender, open operation, Black race, and emergent presentation were independent predictors for a longer stay and greater total charges ( $p < 0.0001$ ). Medicare insurance predicted lower total charges but longer length of stay ( $p < 0.0001$ ). **Conclusion:** Race, insurance, procedure type, and patient presentation influence hospital charges and stays following cholecystectomy. Understanding these trends will allow policymakers and providers to limit the healthcare burden of cholecystectomy.

**Key Words:** Cholecystectomy; Length of Stay; Gallbladder Diseases; Healthcare Costs (Source: MeSH-NLM).

## Introduction

Gallbladder disease is very common in the United States and presents a significant burden to the country's healthcare system. In 2014, cholecystectomy was the 8th most frequent operating room procedure, accounting for 2.6% (372,600) of all operations.<sup>1</sup> Epidemiologists have thoroughly examined identifying and characterizing factors contributing to the high prevalence of gallbladder disease in the United States.<sup>2-5</sup> Variables such as race, gender, and socio-economic class all contribute to the manifestation of this disease.<sup>4,5</sup>

While data on reductions in mortality or morbidity have been conflicting, laparoscopy has been shown to reduce patient hospital stays and total costs.<sup>6-9</sup> First performed in Germany in 1985, the less invasive laparoscopic cholecystectomy quickly became the gold standard approach in the 1990s, replacing the traditional open approach.<sup>10</sup> Despite its now widespread use, healthcare institutions still resort to the traditional open approach under certain circumstances including limited resource settings, lack of qualified surgeons, and predisposing patient risk factors.<sup>11</sup> Whether a procedure is emergent or elective may also determine surgical approach, as some surgeons argue the necrosis and inflammation in acute settings makes laparoscopy unfavorable.<sup>12</sup>

The aim of this study was to identify current trends in cholecystectomy procedures, describe differences in patient characteristics based on surgical approach and admission presentation, and identify factors that predict the patient length of stay and hospital charges. We targeted the

New York State population, as characterization of cholecystectomy procedures in this specific area has not been recently reported.<sup>13</sup>

## Methods

### Patient Population

Patient records were taken from the New York State Department of Health Statewide Planning and Research Cooperative System (SPARCS), a publicly available prospective database that captures all admissions and discharge records from New York State hospitals. All New York State hospitals are required to submit admissions and discharge data, including patient characteristics, treatments, insurance status, and All Patient Refined Diagnosis-related Groups (APR DRG) and International Classification of Diseases, 9th revision, Clinical Modification (ICD-9-CM) codes. The SPARCS database has been used for a variety of clinical and epidemiological studies.<sup>14,15</sup> Due to the open-source nature of the data used for this study, IRB clearance was deemed unnecessary by the research team's affiliated institutional review board.

Admissions and discharge data for 66,647 hospitalizations undergoing non-laparoscopic or laparoscopic cholecystectomies from 2013-2016 were identified using Clinical Classifications Software (CCS) codes for the principal diagnosis of biliary tract disease (149) and the principal procedure of cholecystectomy and common duct exploration (84) (**Table 1**). CCS codes are clustered ICD-9-CM codes that fit into more cohesive and uniform categories, allowing for more effective data analytics. These codes have been used in a variety of clinical studies.<sup>16-18</sup> The data

<sup>1</sup> Bachelor's of Science (BS), California University of Science and Medicine, School of Medicine. Colton, California, United States.

<sup>2</sup> Bachelor's of Art (BA), California University of Science and Medicine, School of Medicine. Colton, California, United States.

**About the Author:** Aria Darbandi is a third-year medical student at the California University of Science and Medicine in Colton, California. His research interests include surgery, surgical outcomes.

### Correspondence:

Christina Chopra

Address: 1501 Violet Street, Colton, CA 92324, US.

Email: [choprac@calmedu.org](mailto:choprac@calmedu.org)

Editor: Francisco J. Bonilla-Escobar

Student Editors: Leah Komer, Duha Shellah, Adnan Mujanovic, Vinson Chan, Vardhmaan Jain, Madeleine Jemima Cox & Nikoleta Tellios

Copyeditor: Sohaib Haseeb

Proofreader: Adam Urback

Layout Editor: Fatma Monib

Submission: Mar 11, 2020

Revisions required: Dec 12, 2020

Received in revised form: May 5, 2021

Acceptance: Jun 9, 2021

Publication: Jul 4, 2021

Process: Peer-reviewed

excluded admissions on non-biliary CCS diagnostic codes such as pancreatic disorder (152) and secondary malignancy (42). This focused our study population to patients with biliary tract disease, whose primary purpose for hospital admissions was a cholecystectomy. Patients under the age of 18 were also excluded. APR DRG codes were then used to create cohorts of patients undergoing laparoscopic (263) and non-laparoscopic (262) procedures.

Primary outcome variables were total hospital charges and length of stay (LOS). Total charges are defined as all hospital expenses accrued from admissions to discharge. LOS is defined as the number of days the patient spends as an inpatient from admission to discharge, rounded to the nearest day. Co-variables used in this study were gender, race, age, presentation, surgical technique, insurance, year of discharge, and APR DRG severity of illness (SOI) score. Patient age was divided into four groups: 18-29, 30-49, 50-69, and above 70. SOI subclasses ranks patients as either minor, moderate, major, or extreme based on the amount of physiologic or organ system function loss. This score was used in our multivariable analysis and considers the severity of secondary diagnosis and comorbidities, accounting for interactions with patient characteristics and requirements for additional resources for care.<sup>19</sup> The insurance category "other" included self-pay, worker compensation, and unreported data. The racial category "other" encompassed multiracial and undisclosed race. Presentation referred to whether the patient was admitted to the emergency department upon admission.

**Statistical Analysis**

Univariable analysis was conducted to summarize total admissions, procedure type (laparoscopic or open), patient presentation (elective or emergency), year of discharge, age group, gender, race, and insurance status. Number of cases and procedure type were described by year of discharge, along with a separate chart outlining yearly changes in LOS and total charges. Comparative analysis was carried out to assess differences between procedure type and patient presentation. Two-sample t-tests were used to compare differences in LOS and total charges among differences in patient presentation and procedure type. Chi-squared tests and two-proportion z-tests were used to compare proportions of the categorical variables of gender, race, age group, insurance status, admissions presentation, and procedure type.

Multivariate linear regression models with selection were used to assess the predictability of outcome variables LOS and total charges. The model included age group, gender, presentation, procedure type, race, insurance status, year of discharge, and SOI score. These factors were chosen because they had significant associations in the bivariate analysis. A two-tailed p-value <0.05 was set for statistical significance for all analyses. All data analysis was conducted using IBM SPSS Statistics 26.0 (Armonk, New York).

**Results**

After this initial screening, 58,141 patient records were included in this study (Figure 1). The characteristics of the study cohort are outlined in Table 2. Admissions for cholecystectomies declined annually, with an overall decrease of 15.3% (15,691 cases in 2013 to 13,602 cases in 2016). The percentage of laparoscopic surgeries increased (91.0% to 92.5%, p<0.0001) (Figure 2). LOS decreased (3.94 to 3.74, p<0.0001) and total charges increased (\$34,260 to \$42,232, p<0.0001) over the course of the study (Figure 3).


Open procedures were more likely to be elective in nature (42.2% vs. 18.6%, p<0.0001) (Table 3). Open procedures were, on average, more expensive and resulted in longer hospital stays (6.88 vs. 3.58 days, \$56,415 vs. \$36,607, p<0.0001) (Table 4). The percentage of emergent presentations increased during the study (78.6% to 81.2%, p<0.0001). Emergent surgeries had longer hospital stays and greater total charges on average (3.97 vs. 3.4 days, \$39,324 vs. \$34,202, p<0.0001).

**Table 1.** Procedural and Diagnostic Codes Included in Study.


Code Type	Code	Description
APR DRG	262	Cholecystectomy except laparoscopic
	263	Laparoscopic cholecystectomy
CCS ICD-9-CM Procedural	84	Cholecystectomy and common duct exploration
CCS ICD-9-CM Diagnostic	149	Biliary tract disease

Legend: APR all patient refined diagnosis-related groups; CCS clinical classifications software


**Figure 1.** Patient Flow Chart



**Figure 2.** Trends in Laparoscopic and Emergent Cholecystectomies



**Figure 3.** Trends in Total Charges and Length of Stay in Cholecystectomies



**Table 2.** Cholecystectomy Patient Demographic and Clinical Characteristics (2013-2016)

Variable	Value
<b>Admissions</b>	<b>58,141</b> 14,532/year
<b>Gender</b>	
Female	37,804 (65%)
Male	30,337 (35%)
<b>Age Group</b>	
18-29	8,333 (14.3%)
30-49	19,251 (33.1%)
50-69	19,731 (33.9%)
70+	10,826 (18.6%)
<b>Race</b>	
White	33,369 (57.4%)
Black	7,407 (12.7%)
Unknown	17,365 (29.9%)
<b>Presentation</b>	
Elective	11,897 (20.6%)
Emergency	46,154 (79.4%)
<b>Operation</b>	
Laparoscopic	53,266 (91.6%)
Open	4,875 (8.4%)
<b>Insurance</b>	
Medicaid	14,977 (25.8%)
Medicare	15,586 (26.8%)
Private	23,248 (40%)
Other	4,330 (7.4%)

**Table 3.** Cholecystectomy Patient Characteristics By Clinical Presentation (2013-2016)

Variable	Emergency	Elective	p value
<b>Mean LOS</b>	3.97± 3.97	3.4± 3.94	<0.0001
<b>Mean Charges</b>	\$39,324± \$33,621	\$34,202±\$38,503	<0.0001
<b>Gender</b>			
Female	30,291 (80.1%)	7,513 (19.9%)	<0.0001
Male	15,863 (78.0%)	4,474 (22.0%)	-
<b>Operation</b>			
Laparoscopic	43,338 (81.4%)	9,928 (18.6%)	<0.0001
Open	2,816 (57.8%)	2,059 (42.2%)	-
<b>Age Group</b>			<0.0001
18-29	7,175 (86.1%)	1,158 (13.9%)	<0.0001
30-49	15,943 (82.8%)	3,308 (17.2%)	<0.0001
50-69	15,092 (76.5%)	4,639 (23.5%)	<0.0001
70+	7,944 (73.4%)	2,882 (26.6%)	-
<b>Race</b>			<0.0001
White	26,209 (78.5%)	7,160 (21.5%)	-
Black	6,011 (81.2%)	1,396 (18.8%)	<0.0001
Other	13,934 (80.2%)	3,431 (19.8%)	<0.0001
<b>Insurance</b>			<0.0001
Medicaid	12,461 (83.2%)	2,516 (16.8%)	-
Medicare	11,505 (73.8%)	4,081 (26.2%)	<0.0001
Private	18,502 (79.6%)	4,746 (20.4%)	<0.0001
Other	3,686 (85.1%)	644 (14.9%)	<0.005

Legend: LOS: Length of hospital stay

Females were more likely to require emergent procedures (80.1% vs. 78.0%,  $p < 0.0001$ ) and underwent laparoscopic procedures more often (93.7% vs. 87.8%,  $p < 0.0001$ ). Black patients were more likely to undergo an emergent procedure than White patients (81.2% vs. 78.5%,  $p < 0.0001$ ). White patients underwent laparoscopic surgeries less often than Black patients (90.9% vs. 91.4%,  $p = 0.10$ ). As the age group increased, the likelihood of emergent presentations decreased (age 18-29: 86.1% vs. 70+: 73.4%,  $p < 0.0001$ ). Proportions of laparoscopic surgeries decreased

**Table 4.** Cholecystectomy Patient Characteristics By Procedure (2013-2016)

Variable	Laparoscopic	Open	p value
<b>Mean LOS</b>	3.58± 3.25	6.88± 6.21	<0.0001
<b>Mean Charges</b>	\$36,607± \$31,051	\$56,415±\$59,197	<0.0001
<b>Gender</b>			
Female	35,414 (93.7%)	2,390 (6.3%)	<0.0001
Male	17,852 (87.8%)	2,485 (12.2%)	-
<b>Presentation</b>			
Emergent	43,338 (93.9%)	2,816 (6.1%)	<0.0001
Elective	9,928 (82.8%)	2,059 (17.2%)	-
<b>Age Group</b>			<0.0001
18-29	8,112 (97.3%)	221 (2.7%)	<0.0001
30-49	18,285 (95.0%)	966 (5.0%)	<0.0001
50-69	17,615 (89.3%)	2116 (10.7%)	<0.0001
70+	9,254 (85.5%)	1,572 (14.5%)	-
<b>Race</b>			<0.0001
White	30,343 (90.9%)	3,026 (9.1%)	-
Black	6,770 (91.4%)	637 (8.6%)	0.10
Other	16,153 (93.0%)	1,212 (7.0%)	<0.0001
<b>Insurance</b>			<0.0001
Medicaid	14,053 (93.8%)	924 (6.2%)	-
Medicare	13,446 (86.3%)	2,140 (13.7%)	<0.0001
Private	21,736 (93.5%)	1,512 (6.5%)	0.0951
Other	4,031 (93.1%)	299 (6.9%)	0.04

Legend: LOS length of stay

as age increased (Ages 18-29: 97.3% vs. Above 70:85.5%,  $p < 0.0001$ ). Medicare patients were the least likely to have emergent operations (73.8% vs. Medicaid: 83.2%,  $p < 0.0001$ ). Medicare patients were also the least likely to undergo a laparoscopic procedure (86.3% vs. Medicaid: 93.8%,  $p < 0.0001$ ).

Multivariable analysis showed that male gender, open procedures, emergent presentation, and Black race predicted significant increases in LOS and total charges ( $p < 0.0001$ ) (Table 5). Medicare predicted decreased hospital charges but longer LOS ( $p < 0.0001$ ).

## Discussion

The findings of this study offer several noteworthy observations. Univariable analysis confirmed the known nature of gallbladder disease. This disease disproportionately affects females, Black people and middle age, generally presents in emergent settings, and is overwhelmingly treated laparoscopically in modern medical practice.<sup>4,5</sup> Interestingly, we found that total admissions have been incrementally decreasing every year in the adult population of New York State, dropping 15.3% from 2013 to 2016. In a New York State study from 1995 to 2013, Alli et al. found that cholecystectomy procedures did not match the increase in population (1.23% procedural increase for a population increase of 6.32%).<sup>13</sup> While the nationwide incidence is rising, we suggest there may be a population-specific fall of all biliary-type diseases in New York State.<sup>4,5,13</sup> Our data did, however, show a rise in emergent admissions, which could be attributed to the specific rise of acute cholecystitis.<sup>2,20</sup> Multi-regional analysis is warranted to better characterize these trends. One possible explanation for this fall in total cholecystectomy procedures is the shifting indications for elective laparoscopic procedures and more thoughtful decision-making by both surgeons and patients, who are better informed about the substantial risks of surgery. We believe this trend will continue in the years moving forward.

In accordance with the literature, we observed a rise in mean total charges and a decrease in LOS over the course of our study.<sup>20</sup> As hospital expenses continue to rise, monitoring ways to limit the economic burden of cholecystectomy becomes more important. We attribute the fall in hospital stays to enhanced patient fast-tracking and the use of multidisciplinary and multimodal teams to expedite rehabilitation.

**Table 5.** Multivariable Predictor for Length of Stay and Hospital Charges

Variables	Length of Stay				Hospital Charges			
	B Coefficient	95% CI		p value	B Coefficient	95% CI		p value
		Lower	Upper			Lower	Upper	
<b>Age</b>								
18-29	-1.04	-1.15	-0.93	<0.0001	-\$7,316	-\$8,426	-\$6,208	<0.0001
30-49	-0.93	-1.03	-0.84	<0.0001	-\$6,114	-\$7,078	-\$5,149	<0.0001
50-69	-0.70	-0.78	-0.61	<0.0001	-\$4,393	-\$5,244	-\$3,542	<0.0001
Over 70	-	-	-	-	-	-	-	-
<b>Race</b>								
White	-	-	-	-	-	-	-	-
Black	.65	.57	.72	<0.0001	\$7,010	\$6,238	\$7,781	<0.0001
Other	.30	.24	.36	<0.0001	\$5,700	\$5,119	\$6,280	<0.0001
<b>Female Gender</b>	-0.08	-0.13	-0.02	0.005	-\$1,605	-\$2,134	-\$1,076	<0.0001
<b>Elective presentation</b>	-1.04	-1.10	-0.98	<0.0001	-\$7,960	-\$8,153	-\$6,748	<0.0001
<b>Laparoscopic technique</b>	-2.35	-2.44	-2.26	<0.0001	-12,766	-\$13,678	-\$11,854	<0.0001
<b>Year</b>								
2013	.34	.27	.40	<0.0001	-\$7,450	-\$8,153	-\$6,748	<0.0001
2014	.19	.12	.26	<0.0001	-\$4,379	-\$5,080	-\$3,678	<0.0001
2015	.05	-0.2	.12	0.186	-\$2,720	-\$3,432	-\$2,010	<0.0001
2016	-	-	-	-	-	-	-	-
<b>Insurance</b>								
Medicaid	.37	.30	.43	<0.0001	-\$1,248	-\$1,908	-\$587	<0.0001
Medicare	.52	.44	.60	<0.0001	\$1,119	-\$317	-\$1,921	0.006
Private	-	-	-	-	-	-	-	-
Other	.15	.06	.25	0.002	-\$4,106	-\$5,087	-\$3,123	<0.0001
<b>SOI Score</b>								
Mild	-1.10	-1.15	-1.04	<0.0001	-\$7,650	-\$8,185	-\$7,114	<0.0001
Moderate	-	-	-	-	-	-	-	-
Major	3.05	2.97	3.14	<0.0001	\$21,203	\$20,327	\$22,079	<0.0001
Extreme	11.15	10.95	11.34	<0.0001	97,660	\$95,630	\$99,690	<0.0001

Comparative analysis between laparoscopic and open procedures suggests that laparoscopy limits hospital costs and patient stay.<sup>3,6,7</sup> Interestingly, open procedures were disproportionately elective in nature. This was noteworthy because there is no indication to prefer the open technique in an elective setting, and some surgeons prefer open procedures in emergent cases due to the associated excess inflammation and necrosis.<sup>12</sup> A 2013 study by To et. al found that conversion rates to open procedures increased nearly two-fold in emergent settings.<sup>21</sup> While the evidence is limited to support using open procedures more frequently in emergent settings, our findings indicate that open procedures are more often used in elective situations.<sup>22</sup> Future research should evaluate the factors that may be influencing this interesting finding.

Geriatric procedures were more often elective in nature and used the open approach. This trend may be explained by concerns that laparoscopy poses increased risk through high physiologic demand, especially considering these patients often present with other comorbidities.<sup>11</sup> For example, insufflating carbon dioxide during laparoscopy may cause acid-base disturbances and changes in cardiopulmonary physiology that are otherwise avoided in the open approach.<sup>23-25</sup> Despite these concerns, systematic studies indicate that laparoscopic procedures in elderly patients offer many advantages, such as lower pain and convalescence, and clinicians still tend to prefer laparoscopy in the elderly in both emergent or elective settings.<sup>11,26,27</sup>

Our multivariable linear regression model illustrated those elective admissions, laparoscopic operations, and younger patient were associated with lower hospital stays and total costs.<sup>3</sup> This model also showed that women had lower total costs and shorter hospital stays than men. Women are more often candidates for cholecystectomies, but men tend to have more complex and longer procedures which may explain their less favorable outcomes.<sup>28,29</sup> In agreement with our findings, Carbonell et al. found in a US-nationwide study in 2000 that male gender was linked to higher charges, longer LOS, and increased morbidity and mortality after cholecystectomy.<sup>3</sup>

Black race was an independent predictor for increased LOS and total cost. Gahagan et al. conducted a 2009-2012 study using nationwide data that had similar findings. Namely, they found that white patients had shorter hospital stays and lower total charges, despite higher morbidity odds.<sup>30</sup> These findings are concerning and warrant further investigation, as they demonstrate a racial disparity in care beyond disease state and presentation. Likewise, compared to Medicaid patients, private insurance predicted a shorter hospital stay, yet a higher total cost. This suggests wealth disparities that could be attributable to several factors including overbilling, or additional treatments and testing. Overall, our data reinforces evidence of racial and insurance-based disparities in healthcare, specifically among cholecystectomy patients.

There are several limitations to this study. The SPARCS database receives administrative coding, which may not be standardized. This could result in variations in coding that alter the assumed specificity of the inclusion criteria used in this study. Additionally, we attempted to limit confounding factors that would influence outcomes by excluding patients with non-biliary primary diagnoses, which means our data does not reflect absolute values of admissions. Comorbidities were also addressed in our multivariable regression model by including APR severity of illness score. While this scoring system is believed to be valid, its efficiency in studies such as ours needs to be further assessed.<sup>19</sup> Furthermore, the SPARCS database accounts for admissions and discharges, meaning an individual patient could account for multiple data entries. Although the SPARCS database has its flaws, it has been used in a variety of epidemiological and outcomes studies and offers great value in assessing trends in the New York State area.<sup>14,15</sup> We suggest caution when inferring these results to nationwide trends. Future studies should include assessing morbidity and mortality, investigating potential causes for disparities seen among specific ethnic groups and insurance types, and examining trends in the pediatric population.

## References

- McDermott K, Freeman WJ, Elixhauser A. Overview of Operating Room Procedures During Inpatient Stays in U.S. Hospitals. Healthcare Cost And Utilization Project. 2014; 2017, 1-18.
- Halpin V. Acute cholecystitis. *BMJ Clin Evid.* 2014 Aug;04-11.
- Carbonell AM, Lincourt AE, Kercher KW, Matthews BD, Cobb WS, Sing RF, Heniford BT. Do patient or hospital demographics predict cholecystectomy outcomes? A nationwide study of 93,578 patients. *Surg Endosc.* 2005 Jun;19, 767-73.
- Stinton LM, Shaffer EA. Epidemiology of gallbladder disease: Cholelithiasis and cancer. *Gut Liver.* 2012 Apr;6(2):172-87.
- Shaffer EA. Gallstone disease: Epidemiology of gallbladder stone disease. *Best Pract Res Clin Gastroenterol.* 2006;20(6):981-96.
- Sandblom G, Videhult P, Crona Guterstam Y, Svenner A, Sadr-Azodi O. Mortality after a cholecystectomy: A population-based study. *HPBA.* 2015;17, 239-243.
- Shea JA, Healey MJ, Berlin JA, Clarke JR, Malet PF, Staroscik RN, et al. Mortality and complications associated with laparoscopic cholecystectomy: A meta-analysis. *Ann Surg.* 1996 Nov;224(5):609-20.
- Fogli L, Boschi S, Patrizi P, Berta RD, Al Sahlani U, Capizzi D, et al. Laparoscopic cholecystectomy without intraoperative cholangiography: Audit of long-term results. *J Laparoendosc Adv Surg Tech A.* 2009 Apr;19(2):191-3.
- Livingston EH, Rege RV. A nationwide study of conversion from laparoscopic to open cholecystectomy. *Am J Surg.* 2004 Sep;188(3):205-11.
- Reynolds W Jr. The first laparoscopic cholecystectomy. *JSL.* Jan-Mar 2001;5(1):89-94.
- Lujan JA, Parrilla P, Robles R, Marin P, Torralba JA, Garcia-Ayllon J. Laparoscopic cholecystectomy vs open cholecystectomy in the treatment of acute cholecystitis: A prospective study. *Arch Surg.* 1998 Feb;133(2):173-5.
- Alli VV, Yang J, Xu J, Bates AT, Pryor AD, Talamini MA, et al. Nineteen-year trends in incidence and indications for laparoscopic cholecystectomy: the NY State experience. *Surg Endosc.* 2017 Apr;31(4):1651-1658.
- Polomsky M, Hu R, Sepesi B, O'Connor M, Qui X, Raymond DP, et al. A population-based analysis of emergent vs. elective hospital admissions for an intrathoracic stomach. *Surg Endosc.* 2010 Jun;24(6):1250-5.
- Bureau of Health Informatics Office of Quality and Health Safety. Statewide Planning and Research Cooperative System (SPARCS). New York State Department of Health.
- Thompson DA, Makary MA, Dorman T, Pronovost PJ. Clinical and economic outcomes of hospital acquired pneumonia in intra-abdominal surgery patients. *Ann Surg.* 2006 Apr;243(4):547-52.
- Chi M Ju, Lee C Yi, Wu S Chong. The prevalence of chronic conditions and medical expenditures of the elderly by chronic condition indicator (CCI). *Arch Gerontol Geriatr.* May-Jun 2011;52(3):284-9.
- Moy E, Coffey RM, Moore BJ, Barrett ML, Hall KK. Length of stay in EDs: Variation across classifications of clinical condition and patient discharge disposition. *Am J Emerg Med.* 2016 Jan;34(1):83-7.
- McCormick PJ, Lin H Mo, Deiner SG, Levin MA. Validation of the All Patient Refined Diagnosis Related Group (APR-DRG) Risk of Mortality and Severity of Illness Modifiers as a Measure of Perioperative Risk. *J Med Syst.* 2018 Mar 22;42(5):81.
- Wadhwa V, Jobanputra Y, Garg SK, Patwardhan S, Mehta D, Sanaka MR. Nationwide trends of hospital admissions for acute cholecystitis in the United States. *Gastroenterol Rep (Oxf).* 2017 Feb;5(1):36-42.
- To KB, Cherry-Bukowicz JR, Englesbe MJ, Terjimanian MN, Shijie C, Campbell Jr. DA, et al. Emergent versus elective cholecystectomy: Conversion rates and outcomes. *Surg Infect (Larchmt).* 2013 Dec;14(6):512-9.
- Andercou O, Olteanu G, Mihaileanu F, Stancu B, Dorin M. Risk factors for acute cholecystitis and for intraoperative complications. *Ann Ital Chir.* 2017;88:318-325.
- Safran DB, Orlando R. Physiologic effects of pneumoperitoneum. *Am J Surg.* 1994 Feb;167(2):281-6.
- Ho HS, Gunther RA, Wolfe BM. Intraperitoneal Carbon Dioxide Insufflation and Cardiopulmonary Functions: Laparoscopic Cholecystectomy in Pigs. *Arc Surg.* 1992 Aug;127(8):928-32; discussion 932-3.
- Hirvonen EA, Nuutinen LS, Kauko M. Ventilatory effects, blood gas changes, and oxygen consumption during laparoscopic hysterectomy. *Anesth Analg.* 1995 May;80(5):961-6.
- Ferrarese AG, Solej M, Enrico S, Falcone A, Catalano S, Pozzi G, et al. Elective and emergency laparoscopic cholecystectomy in the elderly: Our experience. *BMC Surg.* 2013;13 Suppl 2(Suppl 2):S21.
- Musbahi A, Abdulhannan P, Bhatti J, Dhar R, Rao M, Gopinath B. Outcomes and risk factors of cholecystectomy in high risk patients: A case series. *Ann Med Surg (Lond).* 2020 Jan 3;50:35-40.
- Bazoua G, Tilston MP. Male gender impact on the outcome of laparoscopic cholecystectomy. *JSL.* Jan-Mar 2014;18(1):50-4.
- Alqahtani R, Ghnam W, Alqahtani M, Qatomah A, Alkhatami A, Alhashim A. Role of Male Gender in Laparoscopic Cholecystectomy Outcome. *Int J Surg Med.* 2015;1(2):38-42.
- Gahagan JV, Hanna MH, Whealon MD, Maximus S, Phelan MJ, Lekawa M, et al. Racial disparities in access and outcomes of cholecystectomy in the United States. *Am Surg.* 2016 Oct;82(10):921-5.

## Acknowledgments

We would like to thank the New York State Department of Health and SPARCS for allowing open access to the data used in this study. We would also like to thank Jason Crowley, psychometrician at the California University of Science & Medicine, for his assistance in data analysis.

## Conflict of Interest Statement & Funding

The Authors have no funding, financial relationships or conflicts of interest to disclose.

## Author Contributions

Conceptualization: AD, CC. Data Curation: AD, CC. Formal Analysis: AD. Investigation: AD. Methodology: AD. Project Administration: AD, CC. Resources: AD, CC. Software: AD. Supervision: AD, CC. Validation: AD, CC. Visualization: AD. Writing – Original Draft Preparation: AD. Writing – Review & Editing: AD, CC

## Cite as

Darbandi A, Chopra C. Trends and Factors Impacting Healthcare Charges and Length of Stay for Cholecystectomies: A New York State Population-based Analysis. *Int J Med Students.* 2021 Jul-Sep;9(3):202-6.

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

ISSN 2076-6327

This journal is published by the [University Library System](https://www.library.pitt.edu/), [University of Pittsburgh](https://www.library.pitt.edu/) as part of the [Digital Publishing Program](https://www.library.pitt.edu/) and is co-sponsored by the [University of Pittsburgh Press](https://www.library.pitt.edu/).

